

# WINTER MULCH FOR STRAWBERRIES

*Sonia Schloemann, Department of Plant and Soil Sciences, West Experiment Station, University of Massachusetts, Amherst, MA*



An important fall job in commercial strawberry production is mulching. Strawberries are commonly grown in cold climates, such as the northern US and Canada, but the strawberry plant itself is actually quite vulnerable to cold injury. Research has shown that, without mulch, strawberry crowns can suffer damage at temperatures below 12°F and unprotected strawberry plants can suffer desiccation damage from drying winter winds. Protective mulch can protect strawberries from cold by providing insulation, and from desiccation by providing a barrier against drying winds. Mulches will also protect plants from injury caused by soil heaving, which results from freezing/ thawing cycles during the winter. So, a key to consistent quality strawberry production in cold climates is in protecting the plants from severe temperatures or temperature swings through the practice of mulching.

Production systems can also affect the need for mulching. Plants on raised beds, for example, are more vulnerable to cold and desiccation injury than plants in level plantings, especially in locations that are exposed to strong winter winds. Annual production systems, such as fall planted plasticulture, may utilize less hardy or disease susceptible cultivars. As we will see, mulching practices must adapt to these new systems.



**When should the strawberry grower plan to apply mulch?** Research suggests that a good timing guide is to apply mulch after three consecutive days with a soil temperature of 40°F or below. This soil temperature usually occurs after multiple frosts, and when the plants have slowed growth in response to cooler temperatures. It is best to apply mulch before the soil freezes solid. So, in New England, mulches are usually applied in late November.

**What is a good mulch material?** The traditional mulching material for strawberries in New England is straw. Straws from wheat, rice, oats, or Sudan grass work well. Straws coarser than Sudan grass are not recommended. Straw should be clean, free from weed seed, and contains a minimum of

grain seed. Strawberry growers can produce their own straw, often cutting the straw before the grain seed is viable. Store straw for mulching in a dry area. Occasionally, grain seedlings can become a weed problem the following spring; an application of sethoxydim will give good control.

**How much mulch should be applied?** A traditional, level matted row planting will require 2.5 to 3 tons of straw per acre for a 2 to 3 inch deep mulch, or about 300 small bales of average weight. Raised bed plantings and sites with strong wind may require twice this amount for adequate coverage.

**How is the mulch applied?** Smaller plantings may be mulched by hand by shaking out the bales of straw over the row. Larger plantings often use bale choppers to break up the straw bales and distribute the straw over the bed. Choppers are available for both small bales and large round bales.

**How and when is the mulch removed?** In the spring, when plants begin to show growth under the winter mulch (new green tissue), the mulch should be raked off the rows to allow sunlight to penetrate and reach the foliage. Delaying removal will delay plant growth and flowering and may reduce yield. Mulch can be raked off by hand with ordinary yard rakes in smaller plantings. In larger plantings, various mechanical tools are available ranging from modified hay rakes and tedders to equipment specifically designed for the purpose.

**Floating row covers as mulch.** These covers are composed of a plastic such as polypropylene, spun-bonded into a fabric that is permeable to light, air, and water. Research and growers' experiences demonstrate that these covers are useful for winter protection of strawberry plantings. While floating row covers are available in several weights, only the heavier weights are recommended for winter protection. At present a widely available weight recommended for winter strawberry protection is 1.25 oz/yd<sup>2</sup> (42 g/m<sup>2</sup>). A variety of fabric widths are available, with common widths ranging from 15 feet to 60 feet. This material currently costs about 4 cents per square foot. With proper care, this heavier fabric should last 3-4 seasons. Floating row covers are anchored with posts, rocks, or tube sand. The edges may also be covered with soil. Floating row covers are widely used to protect annual plasticulture plantings.

Row covers are best applied on still days. Be sure to line up sufficient labor to place the row cover. If possible, use wider widths for more efficient application. The row cover edges must be anchored, as must areas where 2 covers overlap. A variety of methods are used to anchor the edges. Edges may be anchored with posts, rocks, or tube sand. The edges may also be covered with soil.

Once the mulch is in place, the job is done for the winter. Monitor the planting frequently. If the straw had blown off areas, replace at once. Watch the edges of row covers, and adjust anchors if needed. Repair any rips or holes as possible.

(Reprinted from: New York Berry News, [Vol. 3 No. 11, November 16, 2004](#). Original printing in: *Berry Notes*, Vol. 16 No.1, UMASS Amherst Extension, Nov. 1, 2004)